The Case of the Nighttime Clues

A family guide to bedwetting, sleep, and asking for help.

By Brooke Quinn, MSc Sleep Medicine, RPSGT



READ TOGETHER

Watson has a question:

“Why do some beds get wet while we are asleep?”

Sherlock is ready to help him look for clues.

First, they agree on something important:

A wet bed is not a broken rule.

You are not in trouble. You can tell a trusted grown-up when you need dry clothes, clean bedding, or help. Your body deserves kindness. [1,12]

Our detective job is to notice, wonder, and ask. Grown-ups and health professionals help work out what the clues mean. You do not have to solve this yourself.

For grown-ups: read this page together if it suits your child. Pages 2-4 are for you. The character dialogue is fictional; it does not describe Watson's personal health.

Where sleep fits

Bedwetting can involve development, family history, bladder storage, overnight urine production, and waking to a full bladder. It is unintentional. Sleep apnea is one possible contributor, not a conclusion we can draw from wet nights alone. [1,2]

The sleep connection: disrupted breathing can affect the signals that regulate overnight urine production. Treating confirmed sleep apnea helps some children with bedwetting, but it does not guarantee dry nights. [2,4]

NOTICE | What is happening?

Note wet nights, daytime urgency or accidents, hard or painful stools, and whether wetting restarted after six or more dry months. Bring up frequent wetting or distress with the pediatrician. [1,2]

Mention snoring, mouth breathing, breathing pauses or gasping, repeated waking, morning headaches, and daytime sleepiness or attention/behavior concerns. These are observations to discuss, not a diagnosis. [3]

WONDER | Look beyond a single explanation

Ask about bowel and bladder health as well as sleep. Pediatric guidance uses “enuresis” from age five; younger children are still developing nighttime dryness. [1,2]

ASK | Make a plan together

“My child wets the bed about ___ nights a week. We also notice ___. Could we review bladder, bowel, and sleep factors? What would make a sleep referral appropriate, and when should we follow up?”

Keep support practical and kind

Offer comfortable nighttime protection and waterproof bedding. Encourage actions within the child's control, such as using the bathroom before bed; avoid rewards or penalties based on dry nights. Keep cleanup calm and private. [1] Ask about alarms or prescribed treatment if more help is wanted. [8]

Get help sooner

Prompt medical advice: painful urination, fever, or blood in urine. **Same-day assessment:** new marked thirst and frequent urination, especially with weight loss or unusual tiredness. **Emergency care:** vomiting, belly pain, deep breathing, or confusion with those diabetes symptoms. Do not wait to finish a log. [2,9,10]

Our nighttime case notes

For the grown-up: keep these notes private. This is not a reward chart or a test your child needs to pass. Use what you already know; you do not need to watch all night.

Age: _____ When it started: _____ New or changing? _____

What is hardest for our family? _____

Date	Wet night / daytime symptoms	Sleep or breathing noticed	Bowel symptoms / other notes

A little detective activity - only if your child wants to

NOTICE: “One thing I want my grown-up to know is...”
WONDER: “One question I have about my body is...”
ASK: “The grown-up I can ask for help is...”

Say it, draw it, or write it here. There are no wrong answers.

Leave the appointment with a plan

What are we checking or trying? _____

Follow-up date / who to contact: _____

If questions remain: “What would change the plan or mean we should come back sooner?” This is an original conversation aid, not a validated questionnaire or complete bladder diary.

The science behind the clues

Pediatric guidance includes constipation, daytime urinary symptoms, and upper-airway obstruction among factors to assess in bedwetting. The aim is an individualized evaluation, not a sleep study for every wet bed. [1]

A secondary analysis of a randomized trial included 393 children aged 5-9.9 with nonsevere obstructive sleep apnea. Bedwetting improved more after tonsil and adenoid surgery than with watchful waiting at seven months. A caregiver questionnaire item measured the outcome. Long-term benefit remains uncertain, and this does not make surgery a general bedwetting treatment. [4]

Read more

[1] Canadian Paediatric Society (2023). Evaluation and management of enuresis in the general paediatric setting.

[2] NIDDK. Symptoms & causes of bladder control problems & bedwetting in children.

[3] NHLBI. Sleep apnea in children.

[4] Snow et al. (2021). Enuresis after adenotonsillectomy: secondary analysis of a randomized trial. JAMA Otolaryngology. DOI: 10.1001/jamaoto.2021.2303.

[8] NHS. Bedwetting in children.

[9] NICE. Diabetes in children and young people, recommendations 1.1.1-1.1.2.

[10] NHS. Diabetic ketoacidosis.

[12] NIDDK. Treatment of bladder control problems & bedwetting in children.

About this family guide

Published November 5, 2026. Educational information, not individual diagnosis or treatment. No independent clinical review is claimed. References retain the numbering of the companion article. Links are clickable in the digital PDF.

Watson & Sherlock invite curiosity and offer reassurance. Adults are responsible for arranging care and interpreting symptoms with a qualified health professional. The case notes can stay at home or go to the appointment; they do not need to be shared publicly.